

Table of Contents

Infectious Disease	3	Alcohol Withdrawal	14
Acute Epiglottitis	3	Anemia, including transfusion guidelines....	14
Acute Otitis Media/Sinusitis.....	3	Asthma (Adult & Pediatric).....	15
Anthrax.....	3	Burn Resuscitation.....	15
Cellulitis	4	Diabetes.....	15
Cellulitis, Orbital.....	4	Gastritis/Ulcer/GERD/Dyspepsia	16
Cholera	4	Head Trauma	16
Diarrhea in children & dehydration Rx.....	4	Hypertension	16
Diarrhea, Travelers'	5	Nephrotic Syndrome	16
Fever Definition.....	5	Poisoning	16
Hepatitis	5	Scorpion Bite	16
HIV.....	6	Sickle Cell (Adult & Pediatric)	16
Intestinal Parasites	6	Skin ulcers.....	17
Malaria, including seizure Rx.....	6	Snake Bite	17
Meningitis	8	Transfusion Reaction	18
Necrotizing Fasciitis.....	9	Uterine Prolapse.....	18
Osteomyelitis	9	Pediatrics.....	19
Pneumonia	9	Apnea of newborn or persistent respiratory distress	19
Rabies.....	11	Hypoglycemia	19
Serious Bacterial Infection of the Neonate .	11	Malnutrition	19
Sexually Transmitted Infections	11	Neonatal Feeding	20
Tetanus.....	12	Pediatric Maintenance Fluids	21
Tuberculosis	12	Obstetrics	21
Typhoid Fever.....	12	L&D Culture at BMC	21
Upper respiratory tract infections.....	13	Abruptio of Placenta	21
Urinary Tract Infection	13	Augmentation of Labor	22
Vaginal Candidiasis.....	14	Caesarian Section Orders	22
Adult Medicine.....	14	Chord Prolapse	22
Acute psychosis.....	14		

Chorioamnionitis	22	Trial of Labor After Cesarean (TOLAC):.....	27
Criminal Abortion	23	General Surgery.....	27
Endometritis.....	23	Anesthesia	27
Hyperemesis Gravidarum.....	23	Chest Tube.....	28
Hypertension in Pregnancy	23	Hernia:	28
Malpresentation.....	24	Hydrocele	28
Misoprostol Dosing	25	Postoperative Pain Management.....	28
Postpartum hemorrhage.....	25	Split Thickness Skin Graft.....	28
Molar Pregnancy	25	Wound Vacuum.....	29
Point of Care Ultrasound.....	26	Miscellaneous.....	29
Premature Rupture of Membranes (PROM) 26		Available laboratory/x-ray.....	29
Preterm PROM (PPROM).....	26	Transfer to Tamale	29
Preterm Labor	26	Ward Notes	29

Infectious Disease

Acute Epiglottitis (**H. influenza**)

Chloramphenicol: 25mg/kg Q6hr IV x7d (change to PO when appropriate, max dose 4gm/day)

<1 yr: 6.25 mg/kg PO Q6hr

>1 yr: 12.5-25 mg/kg PO Q6hr

Ceftriaxone: 50 mg/kg IV daily, change to PO cefuroxime at 125mg Q12hr x 10 days

Amox/Clav 1.2g Q6-8hr (7days -12y 30mg/kg q 6-8h)

Acute Otitis Media/Sinusitis

Amoxicillin: Adults: 500-1000 mg Q6-8H x 10 days

Children: 6-12 yrs (<40 kg): 250 mg TID x 10 days

1-5 yrs (<20 kg): 125 mg TID x 10 days

<1 yr: 62.5 mg TID x 10 days

Note: despite above dosing in children <40kg, amoxicillin 80-90 mg/kg/day is preferred over standard dosing due to concentration-dependent resistance of S. pneumonia

Amoxicillin/Clavulanic acid: Adults: 500/125 mg BID -TID x 10 days

Children: 6-12 yrs: 400/57 mg (10 mL) BID x 10 days

1-5 yrs: 200/28.5 mg (5 mL) BID x 10 days

<1 yr: 100/14 mg (2.5 mL) BID x 10 days

Penicillin Allergy:

Co-trimoxazole (sepra)

Adults: 800/160 mg (2 SS tab) BID x 7 days

Children: 6-12 yrs: 400/80 mg (1 SS tab) BID x 7 days

6 months-5 yrs: 200/40 mg (5mL) BID x 7 days

6 wks-5 months: 100/20 mg (2.5 mL) BID x 7 days

Erythromycin - Adults: 250 mg Q6H x 10 days

Children: 2-8 yrs: 250 mg Q6H x 10 days

0-2 yrs: 125 mg Q6H x 10 days

If severe pain or pus still present after 5 days of therapy:

Cefuroxime

Adults: 500 mg BID x 10 days

Children: 250 mg BID x 10 days

Azithromycin

Adults: 500 mg daily x 3 days

Children: 10 mg/kg daily x 3 days

Anthrax: The hospital is in an area endemic for Anthrax, especially the village of Nagbo and the surrounding villages. The presentation is usually cutaneous (an ulcer with eschar and malignant edema) and occasionally systemically presenting with superior vena cava syndrome and severe respiratory distress which is uniformly fatal. One case of anthrax meningitis has been documented. Unlike cases that are related to bioterrorism, the treatment for cutaneous anthrax is Pen VK 500 mg q.i.d x 7 days. Most cases will actually resolve spontaneously over 2-6 weeks. Alternatively, ciprofloxacin 500mg PO bid for 7-10 days can be used. *Severe systemic infections* may be treated

with Penicillin 4-6,000,000 units q.6.h + Cipro 400 IV q8h + Clindamycin 900 IV q8h x 14d. Systemic steroids can also be used, along with drainage of pleural fluid and ascites (Decreases Toxin).

Cellulitis

Outpatient:

Amoxicillin Adult 500-1 g q6 hrs ×7, 6-12 yrs 250 q6 hours ×7, 1-5 years, 125 q6 hrs ×7, <1 yr 62.5 q6 hrs ×7

Flucloxacillin Adult 250-500 q6 hrs ×7, > 10 yrs 250 500 q6 hrs ×7, 2-10 yrs 125-250 q6 hrs ×7, <2 years 62.5-125 q6 hrs ×7

Amox-Clav Adult 500/125 q8 hrs ×7, 6-12 yrs 5 mL 250/62, q8 hrs, 1-6 yrs. 5 mL, 125/30 q8 hrs ×7, 1 mos- 1 yr 0.25 mL/kilogram 125/31 q hrs ×7

Inpatient:

Benzylpenicillin Adult 2 MU IV every 6 hours ×7 children, 250,000/kilogram every 6 hours ×7 or Clindamycin Adult 0.6-2.5 g daily in 3 divided doses, 12-18 yr 150 – 675 mg every 6 hours ×7d, 1 month to 12 yrs 3.75-6.25 mg/KG every 6 hours ×7d

Cellulitis, Orbital: Ceftriaxone adult 2 – 4 gr IV q 24 hrs (children 50 mg/kg q12h) + Metronidazole adult 500mg q8h (children 30mg/kg/dy in 3 divided doses)

Cholera: WHO definition – acute watery diarrhea in a patient 5 years or older with or without vomiting in an area where cholera is likely to occur. Laboratory diagnosis in our situation is impossible. Treatment is primarily fluid management. Antimicrobials are secondary. Doxy 300mg x 1, Tetracycline 500 q6hx3days, Azithromycin 1 g(20mg/Kg in peds) single dose.

Diarrhea in children & dehydration Rx

D/D Acute watery d, cholera, dysentery, persistent d>14 days, d with malnutrition, d associated with recent antibiotic use, Intussusception

The child should be assessed for the degree of dehydration and rehydrated appropriately with either IV solution, intraosseous, or ORS. Antibiotics should not be used routinely. They are only helpful for bloody diarrhea (shigellosis), suspected cholera, and other serious non-intestinal infections such as pneumonia. Anti-diarrheal drugs are useless and dangerous.

Acute Severe dehydration

Infants < 12 months – NS or RL, 6 hours total = 100mL/kg (**30 ml/kg in the first one hour** and then 70ml/kg over the next 5 hours)

12 months to 5 years – NS or RL, 3 hours total = 100ml/kg (**30 ml/kg in the first 30 minutes** and then 70ml/kg over the next 2-1/2 hours)

If unable to place an IV, place an NG tube with ORS solution given at 20mL/kg/hour for 6 hours

If the child is still dehydrated after the above, repeat the IV infusion. If the child is improving the IV can be stopped and ORS given for 4 hours (see below)

Mild to moderate dehydration

Give ORS during the first 4 hours according to the following regimen

< 6kg 200-400 ml

6-9kg 400-700 ml

10-11kg 700-900ml

12-19 kg 900-1400ml or (kg x 75)

This can be given via NGT or the mother can give a teaspoon every 1-2 minutes or frequent sips from a cup.

No dehydration

Breastfeed frequently

ORS – Give mother 2 packets (0-2 years 50 -100mL, >2 years 100- 200 ml, after each loose stool as sips from a cup)

If the child is not exclusively breast-fed, give ORS and food based fluid (soup, rice water, yogurt drinks) or plain water

Continue feeding

Micronutrients – folate, zinc(<6mos 10mg/dyx14, >6mos20mg/dyx10-14 days), vitamin A, iron copper, magnesium

Persistent diarrhea (14 days or longer)

Consider *recurrent viral infections*, non-intestinal infections (pneumonia, sepsis, UTI, oral thrush and otitis media)

If the child has **bloody diarrhea** treat for **Shigella** (ciprofloxacin 10-15 mg/kg BID for 5 days, azithromycin 15mg/kg on day 1, 10mg/kg/dy on day 2-5, or ceftriaxone 50mg/kg to 1.5grams for 2-5 days) Septra and ampicillin are no longer effective. Consider *Intussusception* and amebiasis if no response.

Amebiasis is treated with metronidazole 7.5mg/kg/dose TID for 7-10 days. (Tinidazole 2g qd x 3days)

Giardia is treated with metronidazole 5mg/kg TID for 5 days.

Also consider *intussusception*. **We see it frequently in Ghana.**

Feeding is critical. If a glucose test strip is positive in the stool, give a diet low in lactose such as whole eggs, rice, vegetable oil, glucose and finely ground chicken. Encourage breast-feeding.

Encourage micronutrients as above.

Diarrhea, Travelers’: Caused by enterotoxigenic and enteroaggressive *Escherichia coli*. Only 5% caused by *Salmonella*, *Shigella*, *Campylobacter*, *Giardia*(vomiting and belching), virus or *Amebiasis*. It is only necessary to treat when the patient has more than 3 bowel movements in addition to symptoms (fever, bloody diarrhea). Treatment options are, **Azithromycin 1000 mg as single dose or 500 mg daily x 3 (Children 10 mg/kg daily x3)**, Ciprofloxacin 500mg b.i.d for 1-3 days if sensitive or Rifimaxin 200mg t.i.d. x 3 days. Imodium may be used for symptomatic relief if there is no blood or fever. *Shigella* is covered by Ciprofloxacin 500 bid x 3 days and Azithromycin 500mg day 1 and 250 mg day 2- 5.

Fever Definition: Normal Centigrade 37 (36.1 - 37.2) or Fahrenheit 98.6 (97 – 99)

Each degree C of fever increased over normal = 2 degrees F, i.e. 38C = 100.6 F and 40 C = 104.6 F

Hepatitis: The most common etiology is Hepatitis A, transmitted fecal-orally. Those in the Developing World acquire it in the first few years of life. Hepatitis B & C are primarily transmitted through parenteral, sexual and perinatal exposure. In high prevalence areas HBV is primarily transmitted by vertical exposure from mother to child. It is often asymptomatic with higher rates of chronicity (90%) than infections acquired later in life. In Africa, transmission occurs perinatally and horizontally through ulcerated skin lesions. With hepatitis C, transmission is believed to occur by various percutaneous exposures by injections from conventional and traditional village health workers, dental procedures, cuts from barbers, male and female circumcision. This disease presents itself routinely in the chronic phase. Presentations are often with massive ascites, edema, or GI

bleed (upper or lower). The mainstay of care is primarily symptomatic, as pharmacologic care is not possible. Patients with ascites are often in respiratory distress. Up to 5 liters can be removed at any given time. Diuresis with spironolactone and furosemide is acceptable. A 5:2 ratio of spiro:furo dosing provides the optimum synergistic effect. If portal hypertension is evident, propranolol 10 mg TID may be given to reduce risk of GI bleed. Salt restriction is essential. Always look out for SBP (>250WBC/ml, abdominal pain and fever) which may be treated with Cipro 200 mg IV q12h x 2 days followed by 500 mg po bid x 5 days or Ceftriaxone 2 grams daily x 5 days. E. Coli and Klebsiella are most common agents. Hepatocellular Cancer is common at BMC.

HIV: National Policy refers to Diagnosis in terms of Public Health Number (PH#). 279 is Positive and the 280 is negative. Testing is managed by Chaplaincy. The following Medical Assistant is responsible for coordinating their care: Nelson at Public Health. All cases should be referred to them as they are familiar with the Ghana HIV Protocols.

Intestinal Parasites

Hookworm	Albendazole 400 mgx1 or 200 mg <1yo Mebendazole 500 mg x 1
Cutaneous Larva Migrans	Albendazole 400 mg dailyx7 days
Ascaris	Albendazole 400 mg x 1
Tricuris	Albendazole 400 mg x 1 Mebendazole 500 mg x 1
Pinworms(Threadworms)	Albendazole 400 mg x 1
Esophagostomum	Albendazole 400 mg daily x 5 days
Strongyloides	Albendazole 400 mg q12h x 3 days
H. nana	Praziquantel 25 mg/kg x 1 dose, repeat in 10 days
Giardiasis (Flagellates)	Metronidazole 500mg PO TIDx5d (children 5mg/kg TIDx 5d)
Entamoeba histolytica	Metronidazole 750 mg TID x 10 days (children 7.5-10 mg/kg TID x 5 days)
T. Saginata	Praziquantel 5-10 mg/kg x 1
T. Solium	Praziquantel 5-10 mg/kg x 1
Cysticercosis	treatment with albendazole is controversial due to cerebral reaction to dying parasites
Onchocerciasis	Ivermectin 150 mcg/kg x 1
Bancroftian filariasis	Ivermectin 150 mcg/kg x 1
Shistosomiasis(Mansoni, Haematobium)	Praziquantel 40 mg/kg x1

Malaria, including seizure Rx

The treatment of malaria is now with 2 drugs at all times. Malaria in adults can be treated with Artemether, Artesunate, Amodiaquine, Malarone, Mefloquine, and Quinine in various combinations. The most potent drugs are Artemether related compounds, owing to action on multiple points in the Life Cycle. Quinine should be combined with either fansidar, clindamycin, or doxycycline. Artesunate Combined Therapy (ACT): Artesunate/Lumefantrine and Artesunate (50 mg)/amodiaquine (150 mg base) are the combinations available in Nalerigu.

Please treat only Smear Positive individuals. African Adults, except if pregnant, rarely require IV Quinine.

Inpatient - Severe Malaria

Artesunate Dose <20kg 3mg/kg >20kg 2.4 mg/kg administered IV or IM at the time of admission and then at 12 hours and 24 hours and then daily for a max of 7 days or until taking p.o. . This course is then followed by a full course of ACT x 3d

If parenteral Artesunate is not available, artemether and quinine are acceptable alternatives:

Artemether 3.2 mg/kg IM given on admission and then 1.6 mg/kg at 8hrs and 24hrs and daily until taking PO (up until 4 days). (Adults-160 mg IM followed by 80 mg IM q.d. x4) then follow with ACT x 3d Or Quinine 10 mg/kilogram (max 600mg) q8 hours. If quinine cannot be administered IV, it can be given in the same dosage by IM injection in the anterior thigh. Each dose for IM injection should be diluted in normal saline to a concentration of 60-100 mg salt/mL and injected into 2 sites.

Duration of Parenteral therapy: A minimum of 24 hours in severe malaria.

Complete treatment by getting a full course of Artesunate Combined Therapy (ACT) when patient is able to take oral therapy but not before 24 hours of parenteral therapy

Be vigilant to treat hypoglycemia, anemia, septicemia, pneumonia, seizures, and dehydration. Fluid overload, pulmonary edema, and renal failure are uncommon in childhood malaria in hyperendemic areas.

Gram neg. bacteremia is common (non-typhoidal salmonella) and is treated by Ceftriaxone, Quinolone or Azithromycin.

Hypoglycemia - <5y Glucose < 3 mm/l (<54), >5y Glucose < 2.2 mm/l (<40) - Bolus D10 2.5 ml/kg q15min until normal (Prepare D10 by withdrawing 100ml of fluid from a liter bag of D5 and replacing it with 100 ml of D50.

Maintenance Fluids with Malaria should contain D5 + Isotonic Saline

If ACT is not available or if insurance will not pay, oral quinine may be given as follows in children:

Oral dosing of quinine (50 mg/5 mL) is as follows:

0-3 kg, ½ teaspoon TID

3.1-6 kg, 1 teaspoon TID

6.1-8 kg, 1 ½ teaspoon TID

8.1-11 kg, 2 teaspoon TID

11.1-13.5 kg, 2 ½ teaspoon TID

13.6-16 kg, 3 teaspoon TID

In **Children**, Quinine could be combined with **Clindamycin** 6.7 mg/kg base TID for 7days.

Clindamycin could be also combined with Artesunate 4mg/kg x 3 days or 4mg/kg loading and 2 mg/kg x 6 days without Quinine.

For **adults**, the Quinine dose is 650 mg(salt) TIDx 3-7 days combined with Clindamycin

20mg/kg(max 1.8 grams) of base divided TID for 7 days, or Doxycycline 100mg BIDx7 days.

Outpatient

Artesunate/Amodiaquine

Best post treatment prophylactic effect

Currently in pharmacy, the doses are pre-packaged as Art/Amod (A/A) 100/270 given if greater than 36 kg as one bid x 3 days or if 18- 36 kg one daily x 3 days.

For children (1-8 yo) A/A 50/150 one bid x 3days and for <1yo A/A 1 sachet daily.

You can also write A/A by weight and the pharmacy will figure it out.

Artemether/Lumefantrine **Short Post Treatment Prophylactic effect-important in areas of high transmission. Currently in the pharmacy, the doses are pre-packaged as**

Artemether/Lumefantrine (A/L) (Brand Lumeter) 80/480 one bid x 3 days for adults > 35 kg.

For children 5- 35 kg, dose as follows: 5 - 15 kg A/L 20/120 mg one bid x 3 days, 15-24 kg

A/L 20/120 mg 2 bid x 3days, 24 - 35 kg 20/120 mg three bid x 3 days

Malarone (atovaquone 240/proguanil 100)

Adults and children: 6-8 mg/kg/day for 3 days. In adults, this is equivalent to 4 tab daily x 3 days. This may also be used with artesunate 4 mg/kg/day for 3 days.

Mefloquine

15 mg/kg (base) as an initial dose followed by 10 mg/kg 8-24 hours later. This should be given along with artesunate 4 mg/kg/day.

Anemia with Malaria – Hct 12-15% transfuse 20ml/kg whole blood. If Hgb is 15—21% transfuse if severe condition.

Malaria in Pregnancy

First line for uncomplicated malaria in the 1st trimester is quinine 10 mg/kg TID x 7 days or combined with clindamycin 10 mg/kg/day divided BID (ACT may be used in 1st Trimester if no option). 2nd and 3rd Trimester- use ACT (A/A or AL) – Increased data suggesting that ACT is safe in all trimesters.

Severe Malaria – First line is Artesunate 2.4mg/kg/dy IV/IM on admission, 12, 24 and 48hrs followed by ACT if taking PO and if not may continue Artesunate 4 more day. Quinine IV followed by Oral Quinine + Clindamycin, Alternative in 2nd and 3rd Trimester is Artemether followed by ACT. If Artesunate not available, Artemether preferred over Quinine (Hypoglycemia)

The WHO advises administration of at least three doses of sulfadoxine-pyrimethamine (SP) for IPTp, ideally at each of three antenatal care visits in the second and third trimesters (per WHO guidance, these antenatal visits should occur at 24 to 26 weeks, 32 weeks, and 36 to 38 weeks of gestation) [14,15].

Seasonal Malarial Chemoprevention (SMC)

Pyrimethamine/Sulfadoxine + Amodiaquine qmos during Malaria Season for those 3-59 mos.

Seizures (Ward Protocols are consistent with below)

Diazepam (Dilute 2ml in 8ml saline/glucose)

Newborn-5 yrs: 0.1-0.3 mg/kg IV Q10-15 minutes with a maximum dose of 5 mg daily (may be given rectally if no IV access- dose 0.5mg/kg via cutoff NGT to 3 cm on a syringe)

5-12 yrs: 0.1-0.3 mg/kg IV Q10-15 minutes with a maximum dose of 10 mg daily

>12 yrs: 5-10 mg IV Q10-15 minutes

Phenobarbital

Acute treatment

10-20 mg/kg as initial dose, then 5-10 mg/kg IV Q 15-30 minutes with a maximum dose of 40 mg/kg

Post-seizure

<2 months: 3-5 mg/kg PO/IV daily

>2 months: 5-8 mg/kg/day

WARNING:

The combination of Phenobarbital and Diazepam causes CNS and Respiratory depression.

Prophylactic Phenobarbital is not recommended in Cerebral Malaria

Meningitis – Rapid Timing of 1st Dose is Key to Success

Meningococcal

Ceftriaxone: adult 2-4g IV daily, children 50-80mg/kg/day IV/IM x 10-14d, Neonates 20-50mg/kg once daily for 21d.(Covers Meningo, Pneu, H. Flu) **Be aware that Antibiotics run out so be careful with dosing**

Alternative - Penicillin: Adult 4gMU q4h + Chloramphenicol 1g IV q6h x 14d, Children 200MU/Kg q6h + 25mg/kg q6h x 14d

Dexamethasone 4-10 q6h x 5-7 days for Adults and Children decreases Hearing Loss.(alternative 1-1.5mg/kg/d divided q4-6h (max 16/d) (Benefit not Clear)

Single dose regimens of ceftriaxone and chloramphenicol in oil are effective in epidemic situations

Chemo Prophylaxis – Rifampin 600 bid x 2 days (>1mos 10mg/kg q12h,<1mos 5mg/kg q12h),

Cipro 500 mg x 1 dose, Ceftriaxone 250mg IMx 1(<15yo 125mg IMx 1), Azithro 500mg x 1 (10mg/kg x1)

Practice (Vetted by 2 ID spec) Cipro 500mg on exit + vaccine)

H. Flu Meningitis – Chloramphenicol X 10 days, Ceftriaxone x 10 days, or Ampicillin 200 mg/kg/day x 10 days

Pneumococcal – Penicillin 400,000 units/kg/day q6h x 10 days

TB – fever>14d, fever>7d with family member with TB, Typical CXR, Coma despite Rx, CSF Lymphocytes Give RIPE x 2 mos and a Continuation Phase of INH and Rifampin x 10 mos.

Necrotizing Fasciitis: The infection is usually mixed, so treat for Strep, Clostridium and Anaerobes with antibiotics such as Crystalline Penicillin, Flagyl and Gentamycin IV. *Clindamycin* is very good for Strep. The primary treatment is wide, aggressive surgical debridement. Often more than 2 debridements are necessary. If it is a Buruli ulcer (purple edges that are undermined), wide surgical debridement is the only treatment. Recent evidence shows that a course of Rifampicin with Streptomycin leads to a resolution of small lesions and enables larger lesions to be treated by less extensive excision and reduces the risk of recurrence after surgery. Buruli ulcers are rare in the N. Ghana, as primarily in the South.

Osteomyelitis: This disease is almost always seen in the advanced stages when the affected bone is already dead (the sequestrum) and the new bone (the involucrum) is being formed around it. Antibiotics do nothing for this. Surgery is needed when the involucrum is adequate to replace the affected bone (i.e. adequate for weight bearing on the lower limb). Order an x-ray of the affected limb with 2 views to assess the adequacy of the involucrum. If this is adequate, a sequestrectomy is scheduled. If the involucrum is not adequate, the patient should return in 3 months for follow-up x-ray.

Pneumonia:

Determine PNA score: 1 point per extremes of age (<5, >65), RR (A>30, C>50), P>120, BP <90, Restless and confused, co-morbidities (COPD, CHF, CKD, Multilobes, O2%<92)

OUTPATIENT Pneumonia (Score<2):

No recent antibiotic therapy (>3 months), choose amoxicillin v. Azithromycin v. Cefuroxime:

Amoxicillin

Adults: 1000mg TID x 7d

Children:

50 mg/kg divided BID x 5d

5-12 yrs: 500 mg TID x 7d

1-5 yrs: 250 mg TID x 7d

6 mo–1yr: 125 mg TID x 7d

Azithromycin

Adults: 500mg QD x3-6d

Children: 10mg/kg x3-6d

Cefuroxime

Adults: 500mg PO Q12 x7d

Children:

3mo-12yr: 30mg/kg/day

>12yr: 250-500 q12h x7d

If recent antibiotic therapy (<3 months): use **Amoxicillin/Clav PO**

Adults: 500/125 mg (625 mg) TID x 7 days

Children: 6-12 yrs: 457 mg (10 mL) TID x 7 days

1-5 yrs: 228.5 mg (5 mL) TID x 7

1m – 1 yr.: 114 mg (2.5 mL) TID x 7 days

If penicillin allergy or atypical organism suspected:

Erythromycin

Adults: 500 mg Q6H x 7 days

Children: 8-18 250-500 q6h x 7 days

2-8 yrs: 250 mg Q6H x 7 days

6 months-2 yrs: 125 mg Q6H x 7 days

INPATIENT Pneumonia (Score>2):

Severe – Difficulty Breathing (< 2 mos RR>60 BPM, 2-11 mos RR>50, 1-5yrs RR>40), flaring, retractions, grunting

Amox-clav IV

Adults: 1.2 g IV q8h x 7

Children: 3 mos – 18y 30mg/kg q8h, <3mos 30mg/kg q12h

Ceftriaxone

Adults: 2 g IV daily x 7 days

Children: 50-80 mg/kg daily x 7 days

Ampicillin

Adults: 500 mg IV Q6 or other regimen depending on situation (aspiration)

If aspiration is suspected, add Metronidazole 500 mg IV Q8

Children: 100 -200 mg/kg/day IV/IM divided q6-8h (or benzyl penicillin 200K units/kg/day divided q6h) for at least 3 days and then switch to amoxicillin 25- 50 mg/kg/day divided q12h for 5-7 days if improving.

If doesn't improve but no complications – Cefuroxime 25-30 mg/kg q8h IM/IV.

Consider Staph (pneumatocoeles, empyema) – Flucloxacillin 50mg/kg q6h

IV/IM+Gent

Always consider TB

Adding Doxycycline or Azithromycin would cover Atypical Bact.

Very severe: Cyanosis, lethargy, convulsions, vomiting, not feeding

Adults: Ceftriaxone 2 g qd + gentamycin 2-2.5 mg/kg IV Q8 x 1 week +Doxycycline or Azithromycin.

Children: Ampicillin 200 mg/kg/d div q6-8h (or Ceftriaxone 80mg/kg qd) + Gentamycin 7.5 mg/kg/day x 5 days – if child responds then Amox 15mg/kg po tid +IM Gentamycin x 5 days.

If no response, change to Flucloxacillin 50 mg/kg IM/IV q6h or one may switch Ceftiaxone or Chloramphenicol 75 mg/kg/day divided q8h IM/IV or Cefuroxime 25-50 mg/kg q8h IM/IVx 5 days

Consider bronchodilators, steroids (1mg/kg of prednisolone oral)

Oral therapies:

Amoxicillin/Clavulanic acid: double outpatient doses

NOTE: while effective, the WHO recommends AGAINST use of fluoroquinolones (i.e. ciprofloxacin, levofloxacin) for pneumonia due to role in treating MDR tuberculosis.

Rabies: Uniformly fatal here. Differential diagnosis includes tetanus, drug intoxications, hysteria, Guillain-Barre syndrome, viral encephalitis. For suspected rabies, kill animal and send brain for evaluation to Pong Tamale OR observe animal for 10 days.

Post-bite treatment: immediate cleansing with a virucidal agent and debridement under anesthesia if necessary. Avoid suturing wound. Active immunization for Rabies. (Intradermal regimen below is **no** longer CDC approved) Tissue culture vaccine-0.1 mL intradermal at 2 sites (both deltoids) on days 0, 3, 7, and on day 28 give 0.2 mL at one site (deltoid). A vial of vaccine can remain open in the refrigerator for 8 days, so try to immunize others at the same time. No RIG available. If vaccine plentiful follow label info and **not** Intradermal protocol above. Standard Vaccine at Standard Doses may be available locally or in Tamale. Stop vaccination if the animal survives 10 days.

Post-exposure prophylaxis in previously vaccinated individuals. 0.1 mL intradermal on days 0 and 3, RIG not necessary.

Passive immunization. Rabies Immune Globulin (RIG) 40 IU/kg of equine RIG or 20 IU/kg of human RIG. Inject into wound site and at a remote site (not on buttocks). Not available in Nalerigu.

Serious Bacterial Infection of the Neonate (< 1 mo)

This includes meningitis, pneumonia, and bacteremia

(T>37.5 or <36.5, HR<100 or >160, RR>60 or <20)

Ampicillin 50 mg/kg IV/IM q12h in the first week of life and then q8h between weeks 2 and 4 of life + Gentamicin 4 mg/kg/day given once daily

Or Ampicillin (as above) + Cefotaxime 20-25mg/kg q8h x7 days

Or alternatively Ceftriaxone 100mg/kg/day divided q12h alone (Does not cover Listeria)

Or Cefuroxime IV 25-30 mg/kg q8h (preferred d/t risk of biliary sludge with ceftriaxone)

Omphalitis/Cord Sepsis: Flucloxacillin 15mg/kg IV q6h(<7d and q8h(>7d) + Gentamycin

Treatment Guide: If Term – give IV 3-4 days and if improving change to PO, of if Pre-term(<28wk) give for 7 days after last fever. Good PO regimen is Cefuroxime 30 mg/kg/day in 2 divided doses (Susp. 125/5ml) x 7 days

Remember to treat hypoglycemia, hypoxemia and to keep neonate warm –

See Pediatrics below for hypoglycemia treatment

See Pediatrics below for treatment of apnea of the newborn or persistent respiratory distress

Sexually Transmitted Infections

Chancroid: Ciprofloxacin 500 mg BID x 3 days or Ceftriaxone 250 mg IM

Chlamydia: Doxycycline 100mg PO BIDx7d or Erythromycin 500mg Q6 x7d or Azithro 1g PO x1

Gonorrhea: Ceftriaxone 250 mg IM x 1 or Azithromycin 2 g x 1, will treat both GC and Chlamydia

LGV: Doxycycline 100 mg BID for 14 – 21 days

PID:

Outpatient: Ceftriaxone 250 mg IM + (Doxycycline 100 mg BID X14 days or Azithromycin 1g Q weekly x 2)

Inpatient: Ampillin 2g IV/IM now and 1g IV Q6H + Gentamycin 80 mg IV/IM Q6H + Metronidazole 500 IV Q8H (Metronidazole may be given PV)

Once better, continue for 2 days and then place on Doxycycline to complete 14d of antibiotics.

Syphilis:

Benzathine penicillin 2.4 MU IM x 1 (If latent use 3 doses 1 week apart)(Primary may now require 2 doses)

If penicillin allergic:

Doxycycline 100 mg BID x 14 days
Erythromycin 500 mg Q6H x 14 days
Azithromycin 2 g PO x 1
Ceftriaxone 1 g IM or IV daily x 8-10 days
Plus treat for chancroid as above

Tetanus: Management: prevent toxin release, neutralize unbound toxin, and minimize the effect of already bound toxin, control muscle spasm and autonomic dysregulation

Prompt debridement, quiet room, IV, NGT, Hydration, Feeds (Newborn (NB) breast milk q1h

Adult -Metronidazole 500 mg IV q8h x 7-10 days, Children>1 mos. – Met. 7.5 mg/kg q8h

Penicillin 50,000 unit/kg stat and then 4 MU q6h x5 days is an option but high doses may exacerbate the effects of tetanus (Children 50,000 units/kg q6h x 5 days)

Others – Doxycycline, Clindamycin, Chloramphenicol

If mixed infection, add Ceftriaxone

Neonates – PCN 250,000 units q6h x 7 days + Gentamy4mg/kg IV q24 h x 7 days

Human Tetanus Immune Globulin – Adults and Children 150 units/kg (3000-6000 units), Neonates 500 units IM divided 2 separate sites – see product info for specifics

Complete tetanus immunization as tetanus toxin is poorly immunogenic

Spasms –

Adults – Chlorpromazine 50 mg IM q 4-8 h + Diazepam 10 mg IV or IM q3-6h prn or Phenobarb 200 mg IM q8-12h prn.

Magnesium Sulfate 40 mg/kg over 30 min. followed by 1.5g(<45yo) – 2g(>45yo) per hour to control muscle spasms and autonomic dysfunction. (See Pre-Eclampsia for dosing help)

Labetalol infusion, Morphine Sulfate

Children – Diazepam IV/IM/PR 3-6 mg q3-6h prn + Phenobarbital 5mg/kg IM, NGT stat and then 2.5 mg/kg q 12 h, or Chlorpromazine 12.5-25mg IM, NGT q8h prn

Neonates – Chlorpromazine 7.5 mg IM or NGT q8h + Phenobarbital 30 mg stat and 7.5 mg IM or NGT q 12h, or Diazepam 2 mg IV, IM, NGT q3-6h prn

Tetanus prevention for Deep Wounds is ATS (anti-tetanus toxin) 1500 units IM (no Tetanus Toxoid available)

Tuberculosis: Common presentations are pulmonary, spinal (Pott's disease), TB peritonitis, TB Pericarditis(pericardial effusion= TB) and scrofula. If TB is suspected, a chest x-ray and 2 sputums may be collected (spine films if Pott's is suspected). Sputum may be negative in patients with TB (up to 50%); clinical suspicion should be taken into account, especially in wasting children. RIPE treatment (rifampin, isoniazid, pyrazinamide, ethambutol) is initiated and driven by public health. The special needs department of the hospital handles TB patients. If the patient is sick enough to be admitted, they should be admitted to TB isolation where they get RIPE treatment for 2 weeks before being discharged to the TB village. If they do not need admission, they may be sent to the TB village to start DOT therapy.

Typhoid Fever (Non-focal Fever>7 days without Malaria)

Ciprofloxacin 500 mg PO or 400mg IV q12h x 7-10 days in adults

Children PO 15mg/kg q12h or IV -10mg/kg q12h PO x 7-10 days

Ceftriaxone: Adult: 2-4 g daily x 10-14 days

Children: 50-100 mg/kg IV daily x 10-14 days

Azithromycin: Adult: 500- 1 g PO daily x 5-7 days

Children: 10-20 mg/kg (up to 1 g) daily x 5-7 days

With Ciprofloxacin resistance (SE Asia), this drug will become more important.

In severe cases with depressed LOC or shock, dexamethasone 3 mg/kg IV loading dose followed by 1 mg/kg Q6H x 8 doses of steroids (controversial and the high dose will deplete our Dexa)
Chloramphenicol 2-3 g divided QID or 75-100 mg/kg divided Q6 x 10-14 days
Amoxicillin 100 mg/kg/day divided TID x 2 weeks (resistance)
Septra 8-10/kg/day TMP divided QID (resistance)
Be vigilant to treat children for this with persistent fever and no evidence of malaria. Be cognizant of these patients developing acute abdomens from ileal perforation.

Upper respiratory tract infections

Acute Bronchitis (complicated)

Amox-Clav Adults: 1 g q12h x 5-7 days
Amox Clav Children (>12 y 500mg q12h, 6-12y 5ml(400/57 suspension) q12h x7, 1-6y 2.5ml q12hx7, 1mos-1y 2.5ml(125/31 suspension) q12h
PCN allergy – Azithromycin 500mg daily x 3, Children 10mg/kg daily x 3

Pharyngitis/Tonsillitis (streptococcal)

Amoxicillin

Adults: 500 mg Q6H x 10 days
Children: 6-12 yrs: 250 mg Q6H x 10 days (**Often no fridge in village**)
1-5 yrs: 125 mg Q6H x 10 days
<1 yr: 62.5 mg Q6H x 10 days

Oral Penicillin is 250 tid for 10 days or for a child <12yo 25mg/kg/day x 10 days

Erythromycin (if penicillin allergy)

Adults: 500 mg Q6H x 10 days
Children: 2-8 yrs: 250 mg Q6H x 10 days
0-2 yrs: 125 mg Q6H x 10 days

Urinary Tract Infection

Uncomplicated:

Co-trimoxazole (Bactrim)

Adults: 400/80 mg 2 BID x 3 days
Children: 6-12 yrs: 360-720 mg (7.5-15 mL) BID x 3-5 days
1-5 yrs: 180-360 mg (3.75-7.5 mL) BID x 3-5 days

Ciprofloxacin

Adults: 250 mg BID x 3-5 days
Children: not indicated

Cefuroxime

Adults: > 12yo: 500mg BID x 7 days
2 - 12yo: 15mg/kg(max 250mg) BID x 7 days.
3mos – 2yo: 10mg/kg(max 125) BID x 7days

Complicated:

Ampicillin 1-2 g IV QID PLUS Gentamicin 2.5 mg/kg IV BID x 7-10 days OR

Ciprofloxacin 500 mg BID x 7-10 days

Pyelonephritis: Ceftriaxone 1-2 g IV daily x 14 days

Ciprofloxacin 500 mg IV BID x 14 days

Ampicillin 1-2 g IV QID PLUS Gentamicin 2.5 mg/kg IV BID x 14 days

Change to PO drug when appropriate

In pregnancy, co-trimoxazole (septra, bactrim) can be used in the 2nd trimester. Cipro should be avoided.

Vaginal Candidiasis

Oral: Fluconazole 150 mg PO x 1

Vaginal: Clotrimazole 100 mg x 2 vaginal tabs nightly x 3 days OR

2% cream 5 g vaginally nightly x 3 days OR

Nystatin 100,000 u vaginal tab nightly x 7-14 days

Adult Medicine

Acute psychosis

The only anti-psychotic drug is chlorpromazine (Largactil). A typical dose is 50-100 mg IM followed by 25mg PO b.i.d. Valium is also available for sedation.

Alcohol Withdrawal

Syndrome	Clinical Findings	Onset after last drink
Minor withdrawal	Tremulousness, mild anx, HA, diaphoresis, palpitations, anorexia, GI upset, nml MS	6-36 hrs
Seizures	Generalized tonic-clonic, short postictal	6-48hr
Alcoholic hallucinosis	Visual/aud/tact hallucinations, orientation intact, nml VS	12-48hr
Delirium tremens	Delirium agitation, tachycardia, HTN, fever	48-96hr

- If in active delirium tremens: Phenobarbital 260mg IV once, then 100mg PO Q8hr x6 doses, then 60mg PO TID x6 doses, then 30mg PO TID x6 doses.
- If h/o DT: 100mg PO Q8hr x6 doses, then 60mg PO TID x6 doses, then 30mg PO TID x6 doses.
- If no h/o DT or mild withdrawal: 60mg PO TID x6doses, then 30mg PO TID x6 doses.
- Supplement with diazepam 5mg PO or IV Q4hr PRN agitation
- Fluids: DS NS @maintenance rate. Be mindful that pt may be severely hyponatremic and rapid correction ca be dangerous; switch to PO supplementation ASAP.
- Supplements: daily multivitamin, folate 5mg, thiamine (if available), KCl 600mg tab (can start with KCl IV if not taking PO). Also add Mag sulfate if severe withdrawal.

Anemia, including transfusion guidelines

50% of the anemia is caused by iron deficiency. It normally coexists with a number of other causes including: Malaria, parasitic infections (Hookworm, Trichuris Trichiura, Schistosomiasis), Parvovirus B19, other nutritional deficiencies (folate, B12, vitamin A), hemoglobinopathies (sickle cell disease, thalassemia, hemoglobin C disease), inflammatory conditions causing anemia of chronic disease (HIV), hypersplenism. The highest risk groups are pregnant women and young children.

Definition(hemoglobin): 6 months to 5 years <11, 5 - 11 years<11.5, adult male<13, female<12.

Treatment:

a)Iron supplements - Children (elemental iron 5 mg/kilogram/day), adults (60 mg elemental iron/day) Preparations available: Ferrous sulfate 200mg(65 mg of elemental iron), Iron Polymaltose 3(50mgof elemental iron)

- b) Hookworm, Trichuris - albendazole 400 mg x1(not in first trimester)
- c) Malaria (see previous sections)
- d) Schistosomiasis – Praziquantel(adult-40 mg per kilogram in divided doses x1) (child >4yo 40 mg per kilogram divided doses bid x 1 day)
- e) Folate deficiency - Folic acid 1 mg daily (1-11mos – 15mcg/kg po qd, 1-10yo (1mg daily)
- f) B12 deficiency - B12 1 mg po daily (injectable not available)
- g) Vitamin A – see malnutrition, Pregnancy 2500 units daily, non-pregnant 2300u, male 3000u
- h) Transfusion
 - Decompensated anemia
 - Malaria if Hb<4 or if very ill Hgb<6 (Tendency at BMC to transfuse Hb<6 or Hct<18)
 - Sickle Cell Crisis and Hb<5 or 2 below baseline (goal Hgb 7-8)
 - Active GI Bleed –Keep Hgb>9 until rebleeding unlikely
 - Pregnant < 36 wks. and Hgb<5, >36 wks. and Hgb<6
 - Pregnant <36 wks. with CHF, Malaria, severe infection, heart disease and Hgb<7, >36 wks and Hgb<8.
 - Infants and children and Hgb<4
 - Surgery for minor procedures and Hgb<7-8
 - Major surgery or decompensated patient may require transfusion at higher Hgb

Asthma (Adult & Pediatric)

Acute- emergency

O2 – Nasal 2-6 liters

Nebulization - Salbutamol 2.5 -5mg (may repeat initially after 15 min x 1) then q2h until stabilized (children 2.5 mg or 0.5ml of 5/ml vial.

Hydrocortisone 200mg IV stat, then 100 mg IV q6h. (Children 6-12y 100mg q8h, 1-5y 50 q8h, <1y 5mg q8h)

For severe cases unresponsive to above, aminophylline 250 mg IV(over 20 min) Q6 may be given x 24h. In children the aminophylline loading dose is 3-5 mg/kg (given over 20 min)(max 300mg) and then q6h (Care for Toxicity).

Adrenalin (1:1000) 0.01 ml/kg up to 0.3 ml Q15 minx 2 if salbutamol not available. MgSO4 50% bolus 0.1 ml/kg (50 mg/kg) over 20 min. (MgSO4 not in Ghana Protocol)

Salbutamol oral (1-5 y) child 2mg tid) is available.

Acute –exacerbation

Nebulization – Salbutamol 2.5-5 q6h

Prednisolone Adult 30-40 mg daily (children 1-2 mg/kg/day) PO x 5 days is available.

Dexamethasone (child 0.3mg/kg/dy divided BID) IV but has never been shown more effective than oral steroids.

Aminophylline 200 mg PO QID is available.

Burn Resuscitation for 40-80 kg person – 10ml/hr x TBSA burned (2nd and 3rd) = vol/hr
e.g. 70 kg person with 40% burn – 10 x 40 = 400 ml/hr or 9600/24hrs

Diabetes: Metformin 500mg tab (500-2000 mg divided bid) is available. Glibenclamide 5mg (2.5-5mg qd-bid) and Glimperide(preferred) 2 or 4mg tabs (1-8 mg qd) are the sulfonylureas available. Insulin availability is variable. Regular insulin or NPH are usually what is available, and patients may be sent home with insulin to be kept in a shaded place. Monitoring is with RBS (random blood sugar) or FBS (fasting).

Constipation: Lactulose 10gm/15 ml 15-30 ml qd or bd and Bisacodyl 5-10 mg

Gastritis/Ulcer/GERD/Dyspepsia

Ranitidine, Omeprazole, Metoclopramide, and Magnesium Trisilicate available

H. Pylori – Amox 1gram bid (or 500mg TID), Flagyl 500 BID, Omepr 20 BID x 10-14 days

Head Trauma: Elevate head of bed 30-45 degrees, O2, normocapnia, deepening sedation and analgesia, normovolemia, temperature control, seizure control, Mannitol 0.25-1 gram/kg IV over 15-20 minutes or Manitol 2% 250 mg tidx24 hr, stress ulcer PPx, enteral nutrition, glycemic control. A Glasgow Coma Score (GCS) of 13 or higher correlates with a mild brain injury, 9 to 12 is a moderate injury and 8 or less a severe brain injury.

Best Eye Response (4)

1. No eye opening
2. Eye opening to pain
3. Eye opening to verbal command
4. Eyes open spontaneously

Best Verbal Response (5)

1. No verbal response
2. Incomprehensible sounds
3. Inappropriate words
4. Confused
5. Orientated

Best Motor Response (6)

1. No motor response
2. Extension to pain
3. Flexion to pain
4. Withdrawal from pain
5. Localizing pain
6. Obeys Commands

Hypertension: First line is bendroflumethazide (BDFZ) (thiazide diuretic), 2.5 or 5 mg (no evidence that 5 mg lowers better than 2.5). *Beta-blocker* is atenolol (50-100mg daily divided bid), *ACE/ARB* is lisinopril (5-80 mg qd), losartan (25-100mg qd), the *calcium channel blockers* is nifedipine sr 20 mg qd or bid. furosemide (20-80 mg qd), aldactone (25-100 qd), aldomet (250-2000 mg daily in 2-4 divided doses), and propranolol (40-240 divided bid) are also available.

Nephrotic Syndrome: Commonly seen as a complication of malaria or the treatment of malaria. Confirm with urine specimen. Treatment is with salt restriction (2-4 g/day), furosemide to control edema, and using lisinopril to control hypertension. Steroids are helpful, prednisone 1-2 mg/kg/day for a couple of months until the proteinuria is resolved for a few weeks, then tapering over months. Maximum dose is 60 mg/day.

Poisoning: This is usually the result of kerosene poisoning or DDT poisoning. Observation is all that is needed. Often have had NGT Lavage with charcoal before get to the bedside.

Scorpion Bite: The scorpion bites in this area produce primarily a local effect of severe pain. Rarely is there systemic envenomation. Treatment is with local injection of lidocaine or marcaine. Potent narcotic analgesics are often necessary. There is pethidine (meperidine), if available, 50-100 mg IM Q4-6 hour PRN or in children 1-1.75 mg/kg PO/SC/IM Q3-4 hour PRN. If given IV, dilute prior to use and administer slowly.

Sickle Cell (Adult & Pediatric)

Routine: Penicillin Prophylaxis (<2) 125 bid and 250 bid (>2) up to 5 yrs and later if no Pnueumoccal Vac and Folic Acid 2.5 – 5 mg qd.

Acute Care: Oxygenation, Hydration, Pain Control, Asses for Infection (CBC, BF, UA, CXR, LP), Transfusion for severe and decompensated anemia

Syndromes:

Osteomyelitis–Staph, Salmonella (IV for 1st 2 weeks with therapy to 4-6 wks.)
Hand-Foot Syndrome (<18mos) Dactylitis–painful swelling of hand&feet
Hemolytic Crisis – deep jaundice, hemoglobinuria, acute drop in Hgb
Splenic Sequestration – Abd Pain, Distension, Shock, rapid Hgb drop
Aplastic Crisis – Pallor, weakness
Vaso-Occulsive (Pain) Crisis – Long bones, abdomen, hands/feet dactylitis
Priapism - acute care + aspiration and if fails glans/corpora fistula
Acute Chest Syndrome – cough, chest pain, hypoxia, tachypnea
CVA – HA, focal findings, Seizure - Acute care and keep Hgb>10, control seizures
Transfusion Indications: Hyperhemolysis, Splenic Seques., CVA, Unresponsive Priapism, Chest Syndrome, hypoxia

Skin ulcers:

A chronic problem for many people. Most are *Tropical Ulcers* caused by staph/strep species. Small ones *with cellulitis* may be treated on an outpatient basis with Augmentin (625 mg TID) or Flucloxacillin (500 mg QID) for 2 weeks in conjunction with daily wound dressing using a topical antimicrobial. Antibiotics should not be given chronically. These patients should also be given vitamin c and advised to increase their protein intake. Large ones are often admitted to the isolation ward and provided wound care and antibiotics. Blood sugars should be checked to rule out diabetes. Rapidly progressing wounds may be indicative of HIV. Patients should receive hospital koko for protein supplementation as well as vitamin C daily. Any dead tissue (black or brown, necrotic, tough, not oozing blood) should be debrided under ketamine anesthesia. Once the wound is composed of red, viable tissue (usually after debridement and antibiotics for several days), skin grafting may be attempted. Usually once a graft has been placed, the patient should be kept for 5-7 days for wound care before discharge.

Consider *Buruli Ulcers* (common in S. Ghana) and *Yaws* (Benzathine Penicillin 0.9 gram IM x 1, E-mycin 250-500 mg or Doxycycline 100 or Tetracycline 500 qid x 14 days. Amox 500 qid x 14.)

Wounds are managed on the wards by the Nurses and as outpatients by the Dresser. Those who are being managed on an outpatient basis require a “Dressing Form” with the number of days of dressings and the followup plan, as in See the Doctor in ?#of days.

Snake Bite:

The most common poisonous snake is the carpet viper (*Echis carinatus*). Its venom acts as an anticoagulant. The other common poisonous snake is the spitting cobra. Cobras are generally provoked into attack, whereas carpet vipers are more aggressive by nature. The effect of the bite is assessed by the whole blood clotting time (WBCT). Assume that bites are never dry bites. If the 20WBCT is reported as “no clot”, order ASV (anti-snake venom) x 2 amp. Stat and 1 amp(10ml) for each additional non-clotting WBCT. Clotting times are ordered twice daily (4 AM and 4 PM generally, and on admission). Adequate coagulation is assumed when there are 3 consecutive 20WBCT of < 5 minutes, and the patient may be discharged home. Give ATS 1500 for Tetanus Px.

The bite of the spitting cobra (*Naja Nigricollis*) causes primarily significant skin necrosis that extends to the fascia. The cobra also spits poison into the eye, creating a snake venom ophthalmia, characterized by corneal abrasions. Treatment is with ASV and, in case of ophthalmia, sterile eye irrigation and /or antibiotic eye drops x 1 week to prevent secondary infection.

Transfusion Reaction

- 1. Stop the blood**
- 2. IV normal saline**
- 3. Adult-hydrocortisone 200 mg IV child-hydrocortisone 100 mg IV**
- 4. Promethazine (Phenergan) Adult 25 mg IV**

Uterine Prolapse – Pessary: measure Pubic Symphysis to sacrum + 1-2 cm, bend vertically to insert under pubic bone. Have woman clean and replace daily, washing with soap and water.

Pediatrics

Anemia, see above under Adult Medicine & Infectious Disease (Malaria section) and below under Malnutrition

Apnea of newborn or persistent respiratory distress (last resort): See SBI above under ID. Aminophylline 6 mg/kg loading and 1-2 mg/kg/dose q8h for respiratory distress, try to decrease to q12h after 3 days and daily after 6 days to avoid toxicity and observe effect.

Dehydration, see above under Infectious Disease (Diarrhea in Children section)

Hypoglycemia

Dextrose 10% 4ml/kg bolus and *immediately* followed by maintenance fluids:

Day 1—D10% at rates below

Day 2 and subsequent, D5% ½ Darrow's (NS 0.18% in D10% in Ghana protocols, but BMC Darrow's does contain some glucose). Of note, at BMC nurses must mix D5 together with ½ Darrow's at equal quantities, as the mixture does not exist.

IMPORTANT: FEED PO/NGT (EBM OR FORMULA) WHENEVER POSSIBLE, AT RATES BELOW, AS IVF RATES CANNOT BE REGULATED PRECISELY

	<37 wk	Term
Day 1	60 ml/kg/day	50 ml/kg/day
Day 2	90	70
Day 3	110	90
Day 4	130	120
Day 5	150	120

Malnutrition

Malnutrition is highly prevalent in the area. The Baptist Medical Centre has an extensive program of inpatient and outpatient care. If one wants the Public Health Dept. to assist with the patient on the ward, they can be asked to come by writing "Feeding List" on your hospital order sheet. They can also be asked to come to assist in the outpatient department. Severe Z score < -3, Moderate < -2 but > -3. There is an outpatient malnutrition program in which patients may graduate to, for post-hospital management. Malnourished children may also be referred there from the outpatient department. For those children with Severe Malnutrition, such as Kwashiorkor (Protein-Energy Malnutrition) and Marasmus, hospitalization may be necessary. 2/3 of deaths from acute severe malnutrition occur within the first week of admission. Special care is necessary. The basic principle is, after initial resuscitation, to give high energy foods with increased protein.

See below:

Resuscitation (days 1-7)

Avoid IV therapy due to the tendency to develop fluid overload.

ORS (Give ReSoMol if available) – 5mL /kg every 30 minutes for 2 hours, then 5-10ml/kg hourly for 4-10 hours. When hydrated start milk (F-75 if available) feeding 130mL/kg/day (100 ml/kg for edematous children)

Plumpy Nut is also available – 6 mo. – 59 mo. give 200 kcal/kg/day (sachet = 500 kcal)

Adults – 2000 kcal/day (4 Sachets)

If IV therapy is required, give Ringers' Lactate with 5% dextrose 15ml/kg over one hour and then 10mL/kg/hour over the next 5 hours.

Specific issues

Diarrhea - lactose intolerance may be treated with yogurt or a cereal/oil/sugar, if available. Treat Giardia with Metronidazole 7.5 mg/kg q8h x 7days.

Hypothermia- keep the child warm

Hypoglycemia - Check blood glucose and if low (54mg/dl or 3mm/l – conversion is 18xmm=mg) give 50 ml of 10% glucose solution or sugared water (1 teaspoon sugar in 3 1/2 tablespoons of water) followed by milk feedings. If glucose remains low, repeat glucose solution. If unconscious or convulsing, give 5mL/kg 10% dextrose IV/NGT

Infection – For mildly sick children showing no signs of infection give Amoxicillin (1mos -1yr 125mg PO bid x 10d, 1-5yr 250 bid x 10d, 5-18yr 500 bid x10d. In very sick children, give Ampicillin 50mg/kg IV q6h for 2-3 days and then oral amoxicillin 15mg/kg q8h for 5 days + Gentamicin 7.5mg/kg daily for 7 days. If the child is not responding after 48 hours add Chloramphenicol or Ceftriaxone in meningitis doses. Consider TB and HIV.

Anemia – for Hgb <4, or 4-6 in the very sick give **10ml/kg** of whole blood

Electrolytes and minerals –potassium supplementation along with zinc and magnesium are important(folic 1mg/d, mvit, zinc 2mg/kg/d, copper 0.3mg/kg/d.

Vitamin A – If vitamin A has not been given within the last month, give as follows

<6 months 50,000 units

6-11 months 100,000 units

>12 months 200,000 units

If any **Xerophthalmia** eye signs (corneal ulceration, clouding and necrosis), give above doses on Day 1, 2 and 14.

Antimalarials- give as indicated if +BF or +rapid test

Intestinal parasites – In children over 12 months, give Mebendazole 500mgx1 dose or 100mg bid x 3 days or albendazole one dose

Post-resuscitation - The Public Health Malnutrition Team will continue to be involved with the patient on the ward and then the patient will be discharged to the Feeding Kitchen for daily care up to 6-8 weeks, as necessary

Additional treatments – Multivitamins with Folate should be used for at least 2-3 months. Iron supplements may begin 2 weeks after admission when the child has regained appetite and starts to gain weight.

Neonatal Feeding

<32 wk Gestational Age will not be able to take 100% of PO goal intake of 120ml/kg/d – divide by 8 and give volume q3h (PO initially and rest by NGT as necessary)

Start low at 5 ml per feed and if no vomiting increase by 25ml/kg/d until at goal.

Premies need 22cal/oz milk. This can be provided in the following ways:

1 – EBM - Add ½ tsp of Formula (FM) Powder to 90 ml (3oz) of EBM. (For 24 cal/oz add 1tsp to 90ml EBM)

2 – FM – 1 ½ scoops FM to 90ml water (20cal/oz FM) + ½ tsp of FM Powder =22 cal/oz

Premies don't get Zinc, Phos, or Iron (transferred in 3rd trimester) so if possible formula is ideal for these nutrients, alternatively: human milk fortified Breast milk 20 kcal/oz.

Goal weight gain 15-20 gm/kg/d

Pacifiers are great for practicing feeding

NB – Regular FM = 1 scoop to 60ml (2oz) =20 cal/oz

Pediatric Maintenance Fluids 2 methods of calculation using NS as the preferred solution

1. 1st 10kg 100ml/kg/dy, 2nd 10kg 50ml/kg/dy, Each added kg 25mg/kg, add 10% every degree of fever.
2. <10 kg 4ml/kg/hr, 10-20kg 40ml/kg/hr for the 1st 10 kg plus 2ml/kg/hr for each added Kg > 10, 20-80kg, 60ml/kg/hr for the 1st 20kg then 1ml/kg/hr for each added Kg>20, max 2.5 liters/day

Seizures, see above under Infectious disease (Malaria section)

Diazepam OR Phenobarbital 10mg/kg stat and then 3-5 mg/kg PO/IV daily

Serious Bacterial Infection of the Neonate (< 1 mo), see infectious disease above

Obstetrics

L&D Culture at BMC, notes from D. Smith regarding how OB at BMC differs from in the US

- BMC is a referral site for high risk OB, be prepared for high M&M
- There are significant differences in the approach to HTN in pregnancy, see below for details
- Local L&D culture is to prefer C-sections over IOL, partially given limited FHR monitoring
 - You will likely get push back for attempting IOL for anything but late term, including if doing IOL for indications that are considered standard per ACOG (i.e. pre-eclampsia at term, twins at 38wk, etc.)
- IOL starts in the morning, regardless of the hour that orders were placed (i.e. if you want to start IOL for pre-eclampsia with severe features at 10PM, you must state, “now,” in your orders and talk with the head midwife)
- Women are hospitalized for 6hr s/p NSVD or 3 days s/p CS
- IUFDs are common:
 - If d/t obstructive labors, destructive procedures (ex. Cephalocentesis in setting of hydrocephalous) are performed to allow for NSVD
 - The fetus should remain attached to the placenta for burial, do not cut the umbilical cord
- Even for preterm infants, formula or fortified breastmilk is not well-accepted
- Local providers did not benefit from NRP training and don’t prioritize respiration during neonatal resuscitations; try to help focus efforts on opening airway and bag-mask breathing
- High prevalence of twins, the 2nd born is considered the older because he/she sent the first twin out to assess the situation
- Babies are named for the day of the week they are born until their naming ceremony of DOL#7

Abruption of Placenta

Pathophysiology: Detachment (often partial) of placenta from uterus before delivery

Signs and Sx: Vaginal bleeding (may be occult), uterine tachysystole, uterine tenderness and/or increased tone between contractions, non-reassuring fetal heart, constant abdominal pain

Management:

1. Assess clotting, failure to clot in 7min = coagulopathy
2. Maternal resuscitation efforts as needed: IVF, transfuse fresh whole blood
3. Delivery:

- a. If term, perform IOL if stable or CS if patient is unstable, non-reassuring FHR present, or remote from delivery
- b. If preterm, administer steroids (see dosing below under, “Preterm Labor”)
 - i. If bleeding resolves, patient stable, and FHR is reassuring, monitor for 7 days and then plan for IOL at 37 wk
 - ii. If severe hemorrhage or concern for worsening maternal or fetal status, deliver

Augmentation of Labor with Oxytocin

Oxytocin may be indicated for augmenting labor in multiparous women whose uterus is hypotonic after several vaginal deliveries. Use caution to assess for cephalo-pelvic discordance and avoid in such instances. The drip protocol is as follows:

- Oxytocin 5U injected into 1000 mL RL/NS (or 2.5U/500ml RL/NS)
- Start drip at 8 drops (2mu/min), increase by 8 drops/min every 30 minutes to a maximum rate of 48 drops (12 mu/min), titrating according to the contraction pattern
- If fetal distress occurs, DC the Oxytocin gtt, place patient on her left side, and administer IVF

Caesarian Section Orders

*Only the patient’s spouse or male family member (i.e. not the patient) can consent to a BTL

Pre-OP Orders

NPO and Notify theater of C-Section
 IV NS 2 liters at fast rate
 Urinary Catheter to drainage
 Amox-Clav 1.2 grams IV now
 Hct/HGB, BF now
 Type and Cross 2U of blood (Provisional)

Post-OP Orders

NPO for 6 hrs after CS
 VS Q1h x4 and then Q6h if stable
 IV NS at good rate
 Advance diet as tolerated
 Urinary Catheter to drainage
 Oxytocin 20U in 500 ml of RL over 4 hrs
 Cytotec 800 mcg rectal x 1 dose
 Amox-Clav 1.2 grams Q8h x 24 hrs
 Gentamycin 80mg IV Q8h x 24 hrs
 Amox-Clav 625 mg Bid x 5 days.
 Diclofenac 75 mg IM q12h PRN x72hr
 Paracetamol 1,000mg PO QID PRN
 Hct/HGB, BF in AM

**On POD#1, change antibiotics to Amox-Clav 625mg PO Q12hr x5 days

***Use, “RL” (Ringer’s lactate) rather than, “LR” (which won’t be understood)

Chord Prolapse

Diagnosis: umbilical cord palpated or visualized distal to presenting fetal part s/p ROM

Rx: Trendelenburg, fill bladder, with hand maintain upward pressure on presenting fetal part until stat CS, proceed with NSVD in setting of IUFD

Chorioamnionitis/Intra-Amniontic Infection (IAI)/Triple I:

Diagnosis: Intrapartum fever to 39C (102.2F) once OR intrapartum fever to 38-39C (100.4-102.1F) with 1 of the following: FHR > 160 x10min, maternal WBC > 15, or purulent discharge from the os (ACOG used to also include maternal tachycardia and fundal tenderness in diagnostic criteria)

Rx: Be prepared for PPH, infection predisposes the uterus to atony → labor dystocia & PPH

- First line: Ampicillin 2gm IV Q6hr AND gentamycin 5mg/kg daily

- Reasonable alternative in Ghana: Amox-Clav 1.2 gm IV Q8hr
- Duration of antibiotic: Until delivery if NSVD OR continue for 24hr s/p delivery if CS
- Cover infant for SBI; in the US, labs risk stratify infants so that antibiotics are deferred in some

Criminal Abortion: Self-induced SAB, patient is charged double the cost of treatment for a SAB.

Endometritis

Diagnosis: Fever x2d on postpartum days 2-10, +/- uterine tenderness and foul-smelling discharge

Rx: Targeting polymicrobial infection

- UpToDate: Clindamycin 600mg PO Q6hr + Gentamycin 4.5mg/Kg IM Q24hr until afebrile 24hr
- BMC Protocol has included combination of 3 antibiotics, change to PO once 48hr afebrile
 - 1st Antibiotic (Choose one): Amox-Clav 1.2gm IV Q8hr, Ampicillin 2g IV/IM and then 1 g IV q6h, Chloramphenicol 1g IV/IM q6h, OR Ceftriaxone 1-2 g q24 hrs
 - 2nd Antibiotic: Metranidazole 500 mg IV q8h
 - 3rd Antibiotic: Gentamycin 80 mg IV/IM q8h

Hyperemesis Gravidarum: Diagnosis: 10% weight loss related to nausea/emesis of pregnancy

Rx (Mild): Metoclopramide 10mg po tid prn or Promethazine 25-50 mg tid prn

Rx (Severe): NS alt with D5W, Metoclopramide 5-10 mg q8h prn and Promethazine 25-50 q8h prn

Hypertension in Pregnancy

Blood pressure	<ul style="list-style-type: none"> • Greater than or equal to 140 mm Hg systolic or greater than or equal to 90 mm Hg diastolic on two occasions at least 4 hours apart after 20 weeks of gestation in a woman with a previously normal blood pressure • Greater than or equal to 160 mm Hg systolic or greater than or equal to 110 mm Hg diastolic, hypertension can be confirmed within a short interval (minutes) to facilitate timely antihypertensive therapy
and	
Proteinuria	<ul style="list-style-type: none"> • Greater than or equal to 300 mg per 24-hour urine collection (or this amount extrapolated from a timed collection) or • Protein/creatinine ratio greater than or equal to 0.3* • Dipstick reading of 1+ (used only if other quantitative methods not available)
Or in the absence of proteinuria, new-onset hypertension with the new onset of any of the following:	
Thrombocytopenia	• Platelet count less than 100,000/microliter
Renal insufficiency	• Serum creatinine concentrations greater than 1.1 mg/dL or a doubling of the serum creatinine concentration in the absence of other renal disease
Impaired liver function	• Elevated blood concentrations of liver transaminases to twice normal concentration
Pulmonary edema	
Cerebral or visual symptoms	
* Each measured as mg/dL.	

Diagnosis: Per ACOG 2013 update on Diagnosis & 2019 update on treatment

- Chronic HTN: BP > 140/90 x2 4hr apart prior to 20wk GA
- Gestational HTN: BP > 140/90 x2 4hr apart > 20 wk GA in the absence of proteinuria

- Pre-eclampsia without severe features: BP > 140/90 x2 4hr apart > 20 wk w/proteinuria
- Pre-eclampsia with severe features: BP > 140/90 with severe feature > 20wk GA
 - Severe features: BP > 160/110, Plt < 100, AST or ALT > 2x ULN, Cr 2x BL or 1.1, pulmonary edema, or cerebral/visual symptoms
 - Proteinuria not required for diagnosis if elevated BPs and severe feature
- HELLP: Hemolysis, elevated liver enzymes, and low platelet count
- Eclampsia: Tonic-clonic seizure in the setting of pre-eclampsia

Management (per ACOG):

- Chronic HTN: Only Rx BP if > 160/105, ASA @12wk, serial growth scans, deliver ~38-39wk
 - Meds: Methyl-dpa, Nifedipine (no labetalol); If meds required, consider delivery 37-38 wk
- Gestational HTN: Assess Q wk for pre-eclampsia (proteinuria, severe feature), deliver 37-38wk
- Pre-eclampsia without severe features: Monitor BP, deliver at 37wk
- Pre-eclampsia with severe features: Deliver if > 34 wk, balance of risks/benefits if < 34wk
 - Eclampsia PPx: MgSO4 IM 5g (one 10 mL vial) in each buttock (10g total), followed by 5 g IM Q4 hours until 24hr s/p delivery
 - Hypertensive urgency (if BP > 160/110): nifedipine 10mg Q20-30min (max 30mg) OR hydralazine 5-10mg IV/IM Q20-30min until BP < 160/110
- Eclampsia: BP control as above, oxygen, FHR changes typically transient, attempt NSVD
 - MgSO4: 4-6gm (8-12 mL of 50% soln mixed with 12-18 mL of NS for 20-30mL load over 20min), then 5gm IM Q4hr until 24hr s/p delivery
 - If refractory repeat loading dose of MgSO4, can also give Valium 5-10mg IV PRN

*Although BP is only treated in pregnancy if > 160/105-110, postpartum Rx threshold is 150/100

Differences in approach to HTN in pregnancy/local practice at BMC:

- Local providers are hesitant to treat elevated diastolic BPs (i.e. when systolic BPs are wnl)
- Dx of severe pre-eclampsia is per the 2002 ACOG Practice Bulletin (not with the 2013 updates)
 - Additional criteria include: BP 160/110 6hr apart (rather than on repeat ~15min), 3+ proteinuria, oliguria, RUQ/epigastric pain, IUGR
- Per the 2017 Ghana Treatment Guidelines (by the Ministry of Health):
 - Published guideline has 150/100 as treatment threshold (instead of 160/110 for PEC or 160/105 for cHTN per ACOG) and states risk of fetal distress with lowering BP < 140/90 but this was routinely done in the prenatal clinic at BMIC in March 2019
 - These Ghanaian guidelines don't discuss timing of delivery but it seems local practice is to give anti-hypertensives and wait for spontaneous labor

Malaria in Pregnancy, see above under, "Infectious Disease"; also PPx Qtrimester as above

Malpresentation

External Cephalic Version: ~50% success rate (higher if with spinal or terb), breech/transverse

- Ideal candidate: multip, 37 wk, fetal head not engaged, adequate AFI
- Contraindications: prior CS (relative), abruption, multifetal gestation, ROM, severe HTN
- Procedure:
 1. POCUS: Assess AFI, confirm fetal presentation and lie, assess for fibroids/bicornuate uterus
 2. Relax the uterus: spinal OR salbutamol 1 vial (Terbutaline 0.25 ml) subQ 5-10 min prior to procedure OR MgSO4 5gm IM 30min prior OR Salbutamol PO (equivalent to Albuterol 4 mg) 60min prior
 3. Lubricate abdomen generously

4. Have 1 clinician apply upwards pressure on the presenting fetal part while another clinician flexes the fetal head towards the fetal chest and applies continuous pressure. Every 1-2 min, stop to assess FHR.
5. Following procedure, check FHT Q5min x 30min

Breech: NSVD reasonable if complete or frank breech and clinician experienced with breech NSVD available (endorsed by ACOG since July 2018 following repeat analysis of Term Breech Trial)

- Contraindications: footling breech (d/t cord prolapse risk), macrosomia, inadequate pelvis, fetal head extension
- CAREFUL mnemonic from ALSO: Check dilation/presentation/cord, Await umbilicus (hands off), Rotate arms, Enter for MSV (don't put finger in fetal mouth), Flex head, back Up, Lift baby
- Strategies for head entrapment: Suprapubic pressure, Dührssen's incisions (cutting anterior cervix at midline), symphysotomy, nitroglycerin (100mcg IV)

Misoprostol Dosing per misoprostol.org, regimens updated 2017

Contraindication: prior uterine scar d/t increased risk of uterine rupture (can use if < 28wk GA)

Associated Risks: uterine tachysystole, meconium-stained amniotic fluid

- Common side effects: GI upset (especially if PO or SL), fever, shivering

For cervical ripening at BMC: Initial dose 25ug given in AM and continued Q4hr until 2-3cntx/10min

Incomplete SAB/retained POC: D&C v. MVA are standard of care over expectant mgnt d/t poor f/u

<13 weeks' gestation	13–26 weeks' gestation	>26 weeks' gestation ⁸	Postpartum use
Pregnancy termination^{a,b,1} 800µg sl every 3 hours or pv*/bucc every 3–12 hours (2–3 doses)	Pregnancy termination^{1,5,6} 13–24 weeks: 400µg pv*/sl/bucc every 3 hours ^{a*} 25–26 weeks: 200µg pv*/sl/bucc every 4 hours ^f	Pregnancy termination^{1,5,9} 27–28 weeks: 200µg pv*/sl/bucc every 4 hours ^{f,9} >28 weeks: 100µg pv*/sl/bucc every 6 hours	Postpartum hemorrhage (PPH) prophylaxis^{12,10} 600µg po (x1) or PPH secondary prevention¹¹ (approx. ≥350ml blood loss) 800µg sl (x1)
Missed abortion^{a,2} 800µg pv* every 3 hours (x2) or 600µg sl every 3 hours (x2)	Fetal death^{a,1,5,6} 200µg pv*/sl/bucc every 4–6 hours	Fetal death^{2,9} 27–28 weeks: 100µg pv*/sl/bucc every 4 hours ^f >28 weeks: 25µg pv* every 6 hours or 25µg po every 2 hours ^b	PPH treatment^{k,2,10} 800µg sl (x1)
Incomplete abortion^{a,2,3,4} 600µg po (x1) or 400µg sl (x1) or 400–800µg pv* (x1)	Inevitable abortion^{a,2,3,5,6,7} 200µg pv*/sl/bucc every 6 hours	Induction of labor^{b,2,9} 25µg pv* every 6 hours or 25µg po every 2 hours	
Cervical preparation for surgical abortion^d 400µg sl 1 hour before procedure or pv* 3 hours before procedure	Cervical preparation for surgical abortion^a 13–19 weeks: 400µg pv 3–4 hours before procedure >19 weeks: needs to be combined with other modalities		

Postpartum hemorrhage:

Be prepared (i.e. know risk factors): multifetal gestation, polyhydramnios, fibroids, prolonged labor, chorioamnionitis, known coagulopathy, C-section, operative vaginal delivery, retained POC

Assess underlying etiology: atony (70%), trauma, tissue (retained), thrombin (coagulopathy)

Rx for atony: bimanual massage, misoprostol 800mcg SL, Oxytocin 20U IV in 500mL RL, Bakri balloon, B-Lynch sutures, uterine artery ligation, hysterectomy, maternal resuscitation (IVF, blood)

Molar Pregnancy

Procedure prep: Cervical block with 10ml 1% Lidocaine at 5 & 7 o'clock at cervico-vaginal junction

- If size < 12wk GA: Misoprostol 400mcg SL 1hr before or PV 3hr before procedure
- If size > 12wk GA: Oxytocin 20U IV in 500mL RL at time of procedure

Procedure: Evacuation via suction D&C (often broken) or MVA, cannula size ≈ weeks GA

Post-Op Rx: Azithromycin 1gm x1

Follow-Up: Defer pregnancy for 1yr, avoid OCPs, give LARC or Depo

- Serial pelvic exams: Q2wk x3month, then monthly x3months, then Q6mo for total of 2yr
- Serial pregnancy tests: Q2wk until neg, then Qmonth x6month (should be neg by 8wk)
- Serial CXR: Monthly, then space once pregnancy test is neg

Point of Care Ultrasound

Fetal Biometry:

- CRL: Freeze, Calc, 1st Trimes, CRL, measure, select, set last point for measurement
- BPD: measure outer skull to inner skull; thought to be more accurate than FL at BMC

FHR via M-mode: OB exam, 2D mode, go to M-mode, adjust line to heart, Update, Freeze, Calc, choose FHR, measure between spikes (between 1 peak if iViz/Sonosite or 2 peaks if stationary US – FHR OB mode, 2D mode, go to M-mode, adjust line to heart, Update, Freeze, Calc, choose FHR, measure between spikes – 1 peak.

AFI: Maintain curvilinear probe parallel to floor (not perpendicular to curve of gravid abdomen)

- AFI – Calc – 2nd screen – AFI, Select Pocket – save, measure each pocket scrolling down 1-4
- AFI is wnl if 5-25 or single deepest pocket 2-8




FIGURE 16.9. Anatomic Landmarks for Biparietal Diameter.

Premature Rupture of Membranes (PROM): ROM prior to the onset of labor at 37+ wk GA

Confirm Dx: fluid flowing from os, pooling, ferning, +Nitrazine test; if equivocal, assess AFI

Management: Some Ghanaian (& European) clinicians will discharge home w/expectant mgnt

- IOL with oxytocin per above gtt guidelines (no evidence to support prostaglandins in PROM)
- Monitor for signs of chorioamnionitis: fetal tachycardia, maternal fever, foul-smelling discharge
- Consider PCN for GBS PPx if ROM > 18hr; no universal GBS screening/PPx program in Ghana

Preterm PROM (PPROM): ROM prior to the onset of labor at < 37wk GA

Confirm Dx: As above. Also assess for infection, abruption, fetal heart rate, cord prolapse

Management:

- Expedient delivery if intrauterine infection, abruption, nonreassuring FHR, risk of cord prolapse, or thick meconium
- If 34- 37 wk and stable plan to deliver, although new info suggests that expectant management until 37 weeks is acceptable. There is still benefit from neonatal steroids for fetal lung maturity.
- If < 34 wk and stable: administer latency antibiotics and neonatal steroids
 - Ampicillin 2gm IV Q6hr x48hrs (or Amox/Clav 1.2gm Q8h), then amoxicillin 500mg TID x5d
 - Alternative: Azithromycin 1gm at time of admission and repeat 5 days later
- If <32 wks in active labor, also add Mag Sulfate (4gm IV slowly over 20 min + IM protocol) for neuroprotection
 - Tocolysis: Goal = allow for admin of steroids. Defer if >4cm or evidence of chorioamnionitis
 - Balance of Risks: Maternal complications easier to manage than neonatal

Preterm Labor

Diagnosis: Regular cntx w/cervical change < 37wk GA

Management: Assess for underlying causes (infection, abruption, substance use, multiples, etc.)

- Steroids for fetal lung maturity: Administer from viability to 34 wk, consider 34-37wk
 - Dexamethasone 6mg Q12 IM x48hr
- Tocolytic therapy: Only administer if < 34wk and for 48hr while completing steroids
 - MgSO4: 6 g (12 mL) IV over 4 min, then 5 g IM Q4
 - Nifedipine : load with 30mg PO, if cntx persist at 90min repeat with 10mg. If labor is successfully suppressed, give maintenance dose of 20mg Q6 x24hr then Q8 x24hr
- Neuroprotection: If < 32 wk, Mag Sulfate (4gm IV slowly over 20 min + IM protocol)
- Perivable: Limit of viability at BMC is ~28wk, gather bag mask and oxygen prior to delivery

Trial of Labor After Cesarean (TOLAC):

Appropriate candidate if 1 prior low transverse CS (no h/o classical, T inversion or J extension). Although not an absolute contraindication, less ideal candidate if short inter-pregnancy interval; local practice at BMC is at least 2yr between deliveries to proceed with TOLAC. Likelihood of success is higher if prior NSVD, prior VBAC, normal BMI, and no prior CS for arrest of dilation or arrest of descent. Risk for uterine rupture is 1:200 with risk of serious M&M in 1:2,000. Prostaglandins increase the risk of uterine rupture and are contraindicated. Local practice at BMC is to perform rCS rather than IOL with Oxytocin if a woman who would otherwise be a TOLAC candidate does not go into spontaneous labor by the end of 41wk GA.

General Surgery

Anesthesia

Ketamine (conc. 50mg/ml) 1-2(max) mg/kg IV, may repeat 0.5-1mg after 10 min.

(For IM use-double IV dose – 2.5 ml IV becomes 5ml (max 6mg/kg)

Ketamine causes increased secretions at 8 mg/kg doses, so if using IM anesthesia, must use Atropine which can be mixed with Ketamine. Do not give atropine in a febrile child(hyperthermia)

Premedicate with Atropine(0.015/kg or approx. 0.6 mg vial for an adult) and Diazepam (5-10 mg in adults)

Local: Make Lidocaine with Epi by adding 0.1ml of 1:1000 Epi to 20 ml of 1% Lidocaine

Pressors:

Ephedrine - mix 1 vial (50mg/ml) in 10ml saline and give as a 1ml boluses or 5-10 mg increments.

Adrenaline - mix 1 vial (1/1000 = 1mg) in 1000 ml of fluid or dilute 0.5 mg of Adrenaline(1/2 vial of 1:1000) in 20 ml and give in 1ml increments

Spinal:

Patient/Procedure	Marcaine (0.5%) Dosing	5% Heavy Lidocaine Dosing
Amputation	3 mL	
Hernia	3 mL	2 mL
Prostate	3.5-4 mL	
Adolescent (7yo)	2 mL	1 mL
C-section, small woman	2-2.5 mL	1.2 mL
C-section, medium woman	*May repeat 2.5mL after 8-10min if 1 st spinal fails	1.5 mL (2.5 mL if Lido is 2%)
C-section, large woman		2 mL

Chest Tube

Equipment (when bottles are unavailable): finger of a glove, urinary catheter tubing, chest tube

Procedure: Tie finger of glove to tubing, cut small hole in glove (a one-way valve), connect the tubing to chest tube via a connector, place tube with glove under water in plastic/soda bottle.

Hernia: For a routine inguinal hernia repair, the patient is usually discharged the following day. If it is a complicated hernia, additional days are added depending on the problem. Hydroceles are often kept an extra day.

Pre-Op Orders

Admit to Male/Female Ward
NPO after Midnight
Notify theatre of Hernia Repair in AM
Consent for Hernia Repair signed
Flucloxacillin 1gm IV in AM prior to surgery
(or Amox-Clav 1.2gm IV)

Postop Orders

Bedrest for 6 hours
May eat a regular diet
Pethidine 50mg IM Q4hr PRN pain
(or Diclofenac 75mg IM Q12hr)
Paracetamol 1,000mg PO QID PRN
(Don't write, "Tylenol")

Hernia, Incarcerated: Usually involves inguinal hernias in men. Often these patients may be watched for 1-2 hours before surgery should be alerted. Put the patient in trendelenburg and administer Pethidine (or Diclofenac 75mg IM) and/or diazepam 5-10 mg IV to allow muscles to relax and see if the hernia can be reduced or if it will reduce on its own. If the hernia hasn't reduced and the pain persists, urgent surgical consultation is needed.

Hydrocele

May treat with Sclerosis if high Surgical Risk for Anesthesia (<orange size hydrocoele)
Tetracycline IV solution 500mg -1 gram injected into hydrocoele after Drainage of fluid

Postoperative Pain Management

Adult: Pethidine (Demerol) 50-70mg IM Q4-6hr PRN or Diclofenac 75mg IM Q12hr PRN
Children: Pethidine 1-1.75mg/kg (PO/SC/IM) Q3-4hr

Split Thickness Skin Graft

Equipment: Blades, Knife Handle, Mesher, Saline with Adrenalin, Roll Gauze, ACE, Vasoline Gauze

Procedure:

1. Clean leg with Prep Solution
2. Debride Graft Site and apply adrenalin soaked saline gauze
3. Remove skin with Grafting Handle set at 1 and ½ spaces on scale or by visual
4. Place graft in Saline
5. Apply Gauze impregnated with Saline + Adrenalin to Donor site.
6. Mesh Skin with care to remove skin from the back portion of the Roller Blade. Skin must be attached to Roller Blade to begin process and the skin should be soaked with Saline throughout or it will stick to the Blade.
7. Place the meshed skin on the site and secure it with Staples or Interrupted Sutures.
8. Apply in order: Petrolatum gauze, regular fluffed gauze, roll gauze and then elastic bandage
9. Apply Splint as necessary to control movement
10. Write, "NO DRESSING," on Chart and Dressing itself.
11. First Dressing by Operator on Day #3 and then daily afterwards by ward nurses.

Wound Vacuum: Apply gauze dressing, insert NG tube under gauze, wrap with Saran Wrap and carefully seal the entire dressing and NGT with tape, apply suction for 48 hours.

Miscellaneous

Available laboratory/x-ray

Labs: CBC, blood film (malaria), stool exam, urinalysis, fasting and random blood sugar, sedimentation rate, hepatitis B and C, PH number (HIV), AFB sputum. Cerebrospinal fluid, ascitic fluid, and fluid can be sent to lab for gram stain, WBC with differential, and AFB stain. Electrolytes and CD4 counts are done in Batches on Thursdays (if reagents available).

X-ray machine: Login username = Iddi, password = 123

Transfer to Tamale

Tamale Teaching Hospital (TTH): 037-200-0180 apparently this number is Disconnected and you will have to have a personal contact in the ER to get approval.

Tamale Clinic Schedule: Gen Surg QD, OBGYN QD, Ortho Tues, Urology Fri, Neurology Tues

Ambulance: (Gambaga) 050-601-6616, David (Director) 054-518-2718, Cost C450

National ambulance-193

Ward Notes

Often patients on the wards are followed by a number of physicians. Please write clear, concise notes with good hand writing with a clear diagnosis and plan, even if you are not sure. Long, rambling notes in poor hand writing serve no purpose and compromise patient care. This is especially important when you are leaving and handing over care of your patients to another team.

Phone

Check Minutes *124#

Add Minutes *134*minute number#

“Cure sometimes, treat often, comfort always.” ~Hippocrates